



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : A61K 9/107, 47/10	A1	(11) International Publication Number: WO 94/08564 (43) International Publication Date: 28 April 1994 (28.04.94)
(21) International Application Number: PCT/CA93/00425 (22) International Filing Date: 8 October 1993 (08.10.93) (30) Priority data: 07/957,998 8 October 1992 (08.10.92) US (71)(72) Applicants and Inventors: ALAKHOV, Valery, Yu [RU/CA]; 20726 Gay Cedars, Baie d'Urfe, Quebec H9X 2T4 (CA). KABANOV, Alexander, V. [RU/CA]; 920 Place Sims, #58, Dorval, Quebec H9S 2A1 (CA). SVESHNIKOV, Peter G. [RU/RU]; Teply Stan str., 2-6, Moscow, 117133 (RU). SEVERIN, Eugenie S. [RU/RU]; Novoslobodskaya str., 57/65-65, Moscow, 103055 (RU).		(74) Agent: BLAKE, CASSELS & GRAYDON; P.O. Box 25, Commerce Court West, Suite 2800, Toronto, Ontario M5L 1A9 (CA). (81) Designated States: AU, BB, BG, BR, BY, CA, CZ, FI, HU, JP, KP, KR, KZ, LK, LV, MG, MN, MW, NO, NZ, PL, RO, RU, SD, SK, UA, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: COMPOSITION OF ANTINEOPLASTIC AGENTS INCORPORATED IN MICELLES (57) Abstract Compositions of anti-neoplastic agents incorporate the agent in micelles of at least one block copolymer of poly(oxyethylene)-poly(oxypropylene) in which the ratio of (oxypropylene) blocks to the (oxyethylene) blocks is from about 0.25 to about 1.5 and the micelles have an average diameter of from about 10 to about 25 nm.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NE	Niger
BE	Belgium	GN	Guinea	NL	Netherlands
BF	Burkina Faso	GR	Greece	NO	Norway
BG	Bulgaria	HU	Hungary	NZ	New Zealand
BJ	Benin	IE	Ireland	PL	Poland
BR	Brazil	IT	Italy	PT	Portugal
BY	Belarus	JP	Japan	RO	Romania
CA	Canada	KP	Democratic People's Republic of Korea	RU	Russian Federation
CF	Central African Republic	KR	Republic of Korea	SD	Sudan
CG	Congo	KZ	Kazakhstan	SE	Sweden
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovak Republic
CM	Cameroon	LU	Luxembourg	SN	Senegal
CN	China	LV	Latvia	TD	Chad
CS	Czechoslovakia	MC	Monaco	TG	Togo
CZ	Czech Republic	MG	Madagascar	UA	Ukraine
DE	Germany	ML	Mali	US	United States of America
DK	Denmark	MN	Mongolia	UZ	Uzbekistan
ES	Spain			VN	Viet Nam
FI	Finland				

COMPOSITION OF ANTINEOPLASTIC AGENTS INCORPORATED IN MICELLES

The present invention pertains to improvements in pharmaceutical compositions and in particular improvements in pharmaceutical compositions used in chemotherapy.

5 A number of anti-neoplastic agents currently are in use in chemotherapy (see generally "Cutting's Handbook of Pharmacology, 7th Ed., Chapter 13, Csáky and Barnes) and many additional agents are under investigation.

10 Because of their often complex structure, these agents can exhibit low stability in the blood stream. Many are extremely insoluble and possess poor transport properties with respect to cell membranes. In addition, binding of the anti-neoplastic agent with plasma proteins, as well as other nonspecific interactions in the blood stream prior to its
15 reaching its target, can greatly reduce the effective amount actually available to combat the neoplastic cells. Moreover, multidrug resistance often is observed with such agents; i.e., the sensitivity of the neoplastic cells to the agent is observed to decrease, often by a factor of 10^3 ,
20 over the course of treatment and this resistance thereafter may manifest itself even with respect to structurally different anti-neoplastic agents.

25 In accordance with the present invention, the anti-neoplastic agent of choice (which may include a mixture of several distinct anti-neoplastic agents) is incorporated into a micelle of a block copolymer of poly(oxyethylene)-poly(oxypropylene) in an aqueous dispersion as hereinafter described.

30 The use of the block copolymer micelle in administering the anti-neoplastic agent provides non-covalent solubilization which reduces water-instability and increases the


solubility of the anti-neoplastic agent.

Moreover, while block copolymers of poly(oxyethylene)-poly(oxypropylene) have been used as nonionic surfactants, the effects observed here clearly extend beyond mere solubilization. For example, undesired pretarget protein binding of the anti-neoplastic agent is reduced; i.e., the anti-neoplastic agent appears to be "shielded" from proteins which otherwise would bind to it. Increased sensitivity with respect to the target anti-neoplastic cells also is observed. Finally a reversion in multidrug resistance is observed. While the multidrug resistance (MDR) phenomenon is not fully understood, it is accompanied by an overexpression of a transmembrane P-glycoprotein of M_r about 170 kD (P-170) which mediates the ATP-dependent efflux of numerous drugs from such cells (although drug efflux may involve other membrane components of MDR cells as well). The present compositions appear to possess increased cytotoxic activity with respect to P-170 dependent and P-170 independent MDR cancer cells as compared with sensitive cells, thereby reducing the multidrug resistance effect.

A variety of anti-neoplastic agents are suitable for use in the present composition. These include alkaloids such as vinblastine, colchicine, and demecoline; antibiotics such as those of the rhodomycin group as for example as daunorubicin and doxorubicin, those of the mitomycin group as for example mitomycin C and N-methyl mitomycin C, and those of the bleomycin group such as bleomycin A_2 ; and antifolates such as methotrexate, aminopterin, and dideazatetrahydrofolic acid. It will be appreciated that this improvement extends to mixtures of several such agents.

The present invention is not directed to the underlying anti-neoplastic activity of these agents but rather to an improvement in the manifestation of this activity through formulation.

The block copolymers of poly(oxyethylene)-poly-(oxypropylene) generally are characterized by the structural formula:



I

in which each of x and z , independently of the other, has a value of from about 5 to about 100 and y has a value of from about 20 to about 80. Such block copolymers are known (see Stanton, *Am. Perfumer. Cosmet.* 72(4), 54-58 (1958); Schmolka, *Loc. cit.* 82(7), 25-30 (1967); and *Nonionic Surfactants*, Schick, Ed., (Dekker, NY, 1967) 300-371). A number of these copolymers are commercially available under the generic names of "poloxamers" and "pluronic".

The hydrophobic/hydrophilic properties of a given block copolymer depends upon the ratio of the number of oxypropylene groups to the number of oxyethylene groups. For a composition containing a single block copolymer of poly(oxyethylene)-poly(oxypropylene), for example, this relationship, taking into account the molecular masses of the central hydrophobic block and the terminal hydrophilic blocks, can be expressed as follows:

$$n = \frac{y}{x + z} * 1.32$$

in which y is the number of oxypropylene units and x and z are number of oxyethylene units.

Selecting a block copolymer with the appropriate n value depends upon the hydrophobic/hydrophilic properties of the specific anti-neoplastic agent, or the composite hydrophobic/hydrophilic properties of a mixture of anti-neo-

plastic agents, to be formulated. Typically n will range in value from about 0.25 to about 1.5. This range should be viewed not as numerically critical but as expressing the optimum hydrophobic-hydrophilic balance between the predominantly hydrophilic poly(oxyethylene) blocks and the predominantly hydrophobic poly(oxypropylene) blocks.

An important aspect of the present invention involves utilizing mixture of different block copolymers of poly(oxyethylene)-poly(oxypropylene) to achieve a more specific hydrophobic-hydrophilic balance suitable for a given anti-neoplastic agent or mixture of several anti-neoplastic agents, preserving the optimal size of particles. For example, a first block copolymer may have an n of 1.00 whereas a second may have a value of 1.5. If material having an n of 1.3 is desired, a mixture of one weight portion of the first block copolymer and 1.5 weight portion of the second block copolymer can be employed.

A more generalized relationship therefore for such mixtures can be expressed as follows:

$$N = \frac{y_1 * a}{(x_1 + z_1) * (a + b)} + \frac{y_2 * b}{(x_2 + z_2) * (a + b)} * 1.32$$

in which:

y_1 and y_2 are the number of oxypropylene units in the first and second block copolymers, respectively;
 x_1 and z_1 are number of oxyethylene units in the first block copolymer;
 x_2 and z_2 are number of oxyethylene units in the second block copolymer;
 a is the weight proportion in the first block copolymer; and
 b is the weight proportion in the second block copolymer.

If only one block copolymer of poly(oxyethylene)-poly(oxypropylene) is utilized, N will equal n . An analogous relationship will apply to compositions employing more than two block copolymers of poly(oxyethylene)-poly(oxypropylene).

Using the above parameters, one or more block copolymers of poly(oxyethylene)-poly(oxypropylene) are combined so as to have a value for N of from about 0.25 to about 1.5. The combined copolymers form micelles, the value of N affecting in part the size of the micelles thus produced. Typically the micelles will have an average diameter of from about 10 to about 25 nm., although this range can vary widely. The average diameter of any given preparation can be readily determined by quasielastic light scattering techniques.

The anti-neoplastic compound or compounds in the copolymer micelle are administered parenterally in aqueous formulations, alone or in combination with other therapeutic agents including other anti-neoplastic agents, steroids, etc., to a mammal suffering from neoplasm and in need of treatment. Parenteral routes of administration include intramuscular, intrathecal, intraperitoneal, intravenous and intra-arterial. Isotonic micellar solution of one or more block copolymers of poly(oxyethylene)-poly(oxypropylene) incorporating one or more anti-neoplastic agents are used for parenteral administration. Dosages typically are those associated with the specific anti-neoplastic agent, although as in every case the regimen must be titrated to the particular neoplasm, the condition of the patient, and the response. For example, an isotonic micellar solution of daunorubicin in the block copolymer micelles is administered so as to provide about 1 mg of daunorubicin per kg of body weight. Vinblastine on the other hand is administered in the same fashion but in accordance with conventional usage at lower doses of from about 0.1 to about 0.2 mg/kg. Often

the amount required can be reduced.

The following examples will serve to further typify the nature of the invention but should not be construed as a limitation on the scope thereof which is defined solely by the appended claims.

Example 1

A. A block copolymer of poly(oxyethylene)-poly(oxypropylene) in which $N = 0.25$ (Pluronic F-68) is diluted with RPMI 1640 medium to a final concentration of 2.0% at 4°C. The mixture is incubated for 30 minutes at 37°C and then sterilely filtered through a 0.22 μm filter. An equal volume of a solution of 200 μg daunorubicin in RPMI 1640 medium is added and this mixture is incubated for 30 minutes at 37°C.

B. Human ovarian carcinoma cells (CRL157) are precultured in 1% solution of the same block copolymer but without daunorubicin in RPMI 1640 medium supplemented with 10% calf fetal serum. The preparation of part A is added and the mixture is incubated for 60 minutes at 37°C and the cells then washed three times with RPMI 1640 and cultured in RPMI 1640 supplemented with 10% calf fetal serum for 3 days (Prep. A). Cytotoxicity is measured, both for this preparation and a parallel preparation of free daunorubicin (Prep. B), using the method of Alley et al., *Cancer Res.*, 48, 589-601 (1988). The results are as follows:

conc. (ng/mL)	50000	10000	2000	400	80	16
---------------	-------	-------	------	-----	----	----

% Inhibition

Prep. A	100	100	92	24	6	2
Prep. B	100	81	53	38	20	1

Following the same procedure, cytotoxicity is determined against human T-lymphoma (Jurkat) cells:

conc. (ng/mL)	50000	10000	2000	400	80	16	3.2
---------------	-------	-------	------	-----	----	----	-----

% Inhibition

Prep. A	100	100	100	100	92	33	3
Prep. B	100	100	100	84	51	44	22

Following the same procedure, cytotoxicity is determined against human small cell carcinoma of lung (H-69):

conc. (ng/mL)	50000	10000	2000	400	80	16	3.2
---------------	-------	-------	------	-----	----	----	-----

% Inhibition

Prep. A	100	100	100	100	100	42	12
Prep. B	100	100	100	91	69	42	20

Example 2

Block copolymers of poly(oxyethylene)-poly(oxypropylene) having the ratios of poly(oxypropylene) to poly(oxyethylene) indicated below are dispersed in RPMI 1640 medium at the concentration indicated below. The mixtures are incubated for 40 minutes at 30°C. The average micelle diameter is measured by quasielastic light scattering and the value of *N* calculated as previously indicated. The results are as follows:

	copolymer	conc.	Avg. Diameter	N
5	F-68 ¹	1.0 %	726.0 nm	0.25
	P-85 ²	1.0 %	18.0 nm	1.00
	L-64 ³	1.0 %	20.4 nm	1.50
10	1:1.5 P-85:L-64	0.01 %	17.0 nm	1.30
	1:2.5 F-68:L-64	1.0 %	33.5 nm	1.38

15 Note 1: $x = 80$, $y = 30$, and $z = 80$
 Note 2: $x = 75/2$, $y = 55$, and $z = 75/2$
 Note 3: $x = 27/2$, $y = 30$, and $z = 27/2$

Example 3.

20 A. A 1:1.5 mixture of block copolymers of poly(oxyethylene)-poly(oxypropylene) (Pluronic P-85 and L-64) having individual ratios (n) of (oxypropylene) to (oxyethylene) blocks of 1.00 and 1.50, respectively, and a combined value (N) of 1.30, is diluted with RPMI 1640 medium to a final concentration of 2.0% at 4°C. The mixture is
 25 incubated for 30 minutes at 37°C and then sterilely filtered through a 0.22 μ m filter. An equal volume of a solution of 200 μ g daunorubicin in RPMI 1640 medium is added and this mixture is incubated for 30 minutes at 37°C.

30 B. Cytotoxicity to human ovarian cancer cells (CRL157) is measured, both for this preparation (Prep. A) and a parallel preparation of free daunorubicin (Prep. B) as described in Example 1B. The results are as follows:

conc. (ng/mL)	50000	10000	2000	400	80	16	3.2
---------------	-------	-------	------	-----	----	----	-----

% Inhibition

5

Prep. A	100	100	100	100	94	53	8
Prep. B	100	100	81	50	29	10	2

10

Example 4

Daunorubicin in the composition of Example 3 is evaluated for cytotoxicity in (i) human T-lymphoma (Jurkat) cells as described in Example 1 and (ii) normal human mononuclear cells. The results are as follows:

15

conc. (ng/mL)	50000	10000	2000	400	80	16	3.2
---------------	-------	-------	------	-----	----	----	-----

Cell

% Inhibition

20

Prep. A	Jur.	100	100	100	100	100	74	28
Prep. B	Jur.	100	100	100	83	59	36	21
Prep. A	Norm.	100	100	91	60	21	5	2
Prep. B	Norm.	100	100	80	58	23	18	1

25

Example 5

IC₅₀ values for (i) human T-lymphoma (Jurkat) cells and (ii) normal human mononuclear cells are determined for the daunorubicin composition of Example 3 and compared to those for free daunorubicin. Measurements are made at the indicated intervals of the drug contact with the cells from 15 minutes to 12 hours. The results are as follows:

30

time (hours)		0.25	0.50	0.75	1.0	2.0	4.0	8.0	12
Cell		IC ₅₀ (ng/mL)							
Prep. A	Jur.	150	46	25	17	9.0	0.80	0.50	0.30
Prep. B	Jur.	120	68	35	25	19	16	3.0	5.2
Prep. A	Norm.	3570	950	620	450	250	220	160	140
Prep. B	Norm.	4900	980	405	310	290	275	280	240

Example 6

15 The antineoplastic agent vinblastine is incorporated into the block copolymer mixture described in Example 3. The IC₅₀ of this preparation against SKVO₃ cells, a drug-sensitive human ovarian carcinoma line, is determined to be 0.121 µg/mL; the IC₅₀ against SKVLB cells, an MDR subline
 20 expressing high levels of P-170 obtained through long term cultivation of SKVO₃ in the presence of vinblastine, is 0.0012 µg/mL. The IC₅₀ of free vinblastine against SKVO₃ cells is determined to be 0.095 µg/mL; the IC₅₀ against SKVLB cells is 0.615 µg/mL.

Example 7

25 The antineoplastic agent mitomycin C is incorporated into the block copolymer mixture described in Example 3. The IC₅₀ of this preparation against SKVO₃ cells is determined to be 0.265 µg/mL; the IC₅₀ against SKVLB cells is
 30 0.005 µg/mL. The IC₅₀ of free mitomycin against SKVO₃ cells is determined to be 0.320 µg/mL; the IC₅₀ against SKVLB cells is 1.120 µg/mL.

Example 8

The antineoplastic agent methotrexate is incorporated into the block copolymer mixture described in Example 3. The IC₅₀ of this preparation against SKVO₃ cells is determined to be 0.880 µg/mL; the IC₅₀ against SKVLB cells is 0.0175 µg/mL. The IC₅₀ of free methotrexate against SKVO₃ cells is determined to be 1.090 µg/mL; the IC₅₀ against SKVLB cells is 1.340 µg/mL.

Example 9

The antineoplastic agent colchicine is incorporated into the block copolymer mixture described in Example 3. The IC₅₀ of this preparation against SKVO₃ cells is determined to be 0.720 µg/mL; the IC₅₀ against SKVLB cells is 0.045 µg/mL. The IC₅₀ of free colchicine against SKVO₃ cells is determined to be 0.950 µg/mL; the IC₅₀ against SKVLB cells is 7.450 µg/mL.

Example 10

The antineoplastic agent daunorubicin is incorporated into the block copolymer mixture described in Example 3. The IC₅₀ of this preparation against SKVO₃ cells is determined to be 0.600 µg/mL; the IC₅₀ against SKVLB cells is 0.0068 µg/mL. The IC₅₀ of free daunorubicin against SKVO₃ cells is determined to be 0.620 µg/mL; the IC₅₀ against SKVLB cells is 5.850 µg/mL.

Example 11

To 30 µL of a 20 mg/mL solution of bovine serum albumin in phosphate buffered saline are added 30 µL of daunorubicin

solution in the block copolymer mixture described in Example 3 (Prep. A). A second formulation (Prep. B) is prepared in parallel fashion using free daunorubicin.

5 The preparations are incubated for 10 minutes at 25°C, and then analyzed by HPLC on a TSK-3000 SW gel-filtration column in PBS containing 0.3 M sodium chloride and 5% acetonitrile. Detection is performed at 280 nm and 470 nm. The portion of the drug bound with BSA is determined as:

$$D_b = S_b/S_f$$

10 in which:

S_b is relative area of the 470 nm peak (corresponding to daunorubicin) which coincides in retention time for the 280 nm peak (corresponding to BSA); and

15 S_f is relative area of the peak (or peaks) corresponding to daunorubicin which does not coincide in retention time of the BSA peak.

The results are as follows:

20

Composition	D_b
Prep. A	0.01
Prep. B	0.39

25

Example 12

Micellar daunorubicin obtained as described in Example 3 (Prep. A) and free daunorubicin (Prep. B) are incubated in the dark at 37°C and cytotoxicity to CRL157 cells is then
30 determined in the manner discussed in Part B of Example 1.

The results are as follows:

time (hours)	2	4	12	24	48	96
--------------	---	---	----	----	----	----

5

IC₅₀, $\mu\text{g/mL}$

Prep. A	9.1	10.05	9.8	10.4	10.7	11.3
Prep. B	400	475	1120	6300	10180	48900

10

Example 13

The daunorubicin composition of Example 3 (Prep. A) is evaluated against daunorubicin-sensitive human breast cancer (MCF-7) and two cell lines demonstrating resistance: daunorubicin/verapamil-resistant (MCF-7AU) not expressing P-170, and daunorubicin-resistant, verapamil-sensitive (Dox-MCF-7), expressing P-170, in each case in comparison to free daunorubicin (Prep. B). The results are as follows:

20

conc. (ng/mL)	50000	10000	2000	400	80	16
---------------	-------	-------	------	-----	----	----

Cell % Inhibition

25

30

35

Prep. A	MCF-7	100	100	84	65	42	12
	MCF-7AU	100	100	100	96	69	39
	Dox-MCF-7	100	100	100	89	73	45
Prep. B	MCF-7	100	100	91	69	43	15
	MCF-7AU	100	89	65	37	9	3
	Dox-MCF-7	100	86	62	39	7	2

Free daunorubicin (Prep. B) exhibits higher IC₅₀'s (is less toxic) against both resistant lines. Daunorubicin incorporated in the block copolymers (Prep. A) exhibited lower IC₅₀'s (is more toxic) against both resistant lines.

Example 14

Groups (6 animals/dose point) of C57B1/6 7-week-old female mice are inoculated i.p. with free or micellar (N = 1.3) daunorubicin obtained as described in Example 3 (Prep. B and Prep. A, respectively), and are observed for 14 days. Drug concentrations are adjusted so that a maximum volume of 0.5 mL is injected in each mouse.

The MTD is defined as a dose which leads to no daunorubicin-deaths (any higher dose leads to the daunorubicin-related death of at least 1 animal per group). The experiment is repeated twice. The results are reproducible with less than 10% variation.

The MTD of free and micellar (N = 1.3) daunorubicin is determined to be 2.0 and 1.0 $\mu\text{g/kg}$ body weight, respectively.

Example 15

Daunorubicin possesses high specificity with respect to bone marrow, manifesting itself as reversible leukopenia, i.e., a decrease in the number of WBS (leukocyte count) during drug administration. Bone marrow suppression, as well as anticancer effects of daunorubicin, are conditioned by DNA-intercollating activity, whereas the most harmful side effect of anthracyclines, cardiotoxicity, results mainly from membrane interactions with metabolites (which have low anticancer activity and do not produce significant effects on bone marrow). Therefore, the leukocyte count during *in vivo* administration of MTD daunorubicin allows the assessment of the ratio between specific (DNA-intercollation) activity of the drug and non-specific toxicity.

Groups (6 animals/group) of C57B1/6 7-week-old female mice are inoculated i.p. with free or micellar (N = 1.3)

daunorubicin obtained as described in Example 3 (Prep. B and Prep. A, respectively). Drug concentrations (MTD) are adjusted so that a maximum volume of 0.5 mL is injected in each mouse. Blood samples are collected and viable leukocytes are counted as described in Michisch et al., *Proc. Natl. Acad. Sci. USA* 88, 547-551 (1991). The number of WBC after administration of 0.1 mL PBS, 15-16 mln cells/mL, is used as the control. The experiment is repeated twice. The results are reproducible with less than 10% variation.

The results obtained are as follows:

Days	0	3	7	10	14
------	---	---	---	----	----

WBS, % of control

Prep. A	100	20	46.6	86.6	100
Prep. B	100	40	60	93.8	100

Example 16

The effects of free and micellar daunorubicin obtained as described in Example 3 (Prep. B and Prep. A, respectively) on leukocyte count are determined three days after administration as described in Example 15.

The results obtained are as follows:

Dose of daunorubicin % of MTD	25	50	75	100
----------------------------------	----	----	----	-----

WBS, % of control

Prep. A	85	73	45	21
Prep. B	78	61	36	39

The data shown in Examples 14 through 16 indicate that solubilization of daunorubicin in the block copolymer

micelles does not essentially affect the drug's overall toxicity (MTD of 2 mg/kg and 1 mg/kg for free and micellar drug, respectively), whereas an increase in reversible bone marrow suppression is observed which does not markedly influence the animal survivability.

Example 17

Anti-neoplastic activity is determined by evaluation of the cytotoxic activity of plasma of mammals inoculated with the test composition (see de Valeriola et al., *Cancer Chemother. Pharmacol.* 29, 133-140, 1991).

Groups (6 animals/group) of C57B1/6 7-week-old female mice are inoculated i.v. (via the tail vein) with free or micellar (N = 1.3) daunorubicin obtained as described in Example 3 (Prep. B and Prep. A, respectively). Drug concentrations (MTD) are adjusted so that a maximum volume of 0.1 mL is injected in each mouse. The experiment is repeated twice. The results are reproducible with less than 10 % variation.

To obtain plasma samples, blood (10 μ l) is collected from the tail artery one hour after drug administration, diluted 1:10 with sterile RPMI 1640 medium, and centrifuged at 400 g for 15 minutes. The supernatants obtained are diluted as shown in the table with plasma analogously obtained from mice not inoculated with the drug (the plasma of mice not inoculated with the drug does not produce any significant cytotoxic effect on H-69 cells) and mixed with an equal volume of H-69 cell suspension on RPMI 1640 medium supplemented with 10% fetal calf serum. The cells are incubated for two hours at 37°C and 5% CO₂, and then washed three times with RPMI 1640. The pretreated cells are incubated in RPMI 1640 supplemented with 10% fetal calf serum at 37°C and 5% CO₂ for three days, after which cytotoxicity is determined as described in Example 1.

The results obtained are as follows:

Dilution of plasma	1:20	1:200	1:2000	1:20000
--------------------	------	-------	--------	---------

5

Inhibition, %

Prep. A	100	58	8	0
Prep. B	42	5	0	0

10

Thus cytotoxic titers, the dilution at which the plasma of mice inoculated with preparations B or A produced 50% inhibition of H-69 cell growth, of plasma of mice inoculated with preparations B and A with respect to H-69 cells are determined to be 1:228 and 1:48, respectively.

15

Example 18

The procedure of Example 16 is repeated utilizing SKVLB and SRVO₃ cells. The results are as follows:

a) when MTD of daunorubicin is introduced

5

Plasma Dilution	1:20	1:200	1:2000
-----------------	------	-------	--------

Inhibition, %

10

Prep. A	SKVLB	82	61	18
Prep. B	SKVLB	0	0	0
Prep. A	SKVO ₃	11	0	0
Prep. B	SRVO ₃	9	0	0

15

b) when 10 mg/kg daunorubicin are introduced

20

Plasma Dilution	1:20	1:200	1:2000
-----------------	------	-------	--------

Inhibition, %

25

Prep. A	SKVLB	100	94	69
Prep. B	SKVLB	8	0	0
Prep. A	SKVO ₃	62	31	0
Prep. B	SRVO ₃	22	6	0

30

Example 19

A composition suitable for parental administration is prepared by dissolving 400 mg of Pluronic P-85 and 600 mg of Pluronic L-64 in 50 mL of RPMI 1640 at 4°C. The mixture is incubated for 30 minutes at 37°C and then sterilely filtered through a 0.22 µm filter. This is mixed with a solution of 10 mg of sterile lyophilized daunorubicin powder dissolved in 50 mL of RPMI and incubated for 30 minutes at 37°C.

40

The composition can be stored in the dark at room temperature for 7 days without any essential loss of activity or can be lyophilized and stored for at least 1 year in the dark at room temperature.

Example 20

- 5 A further composition suitable for parental administration is prepared by dissolving 400 mg of Pluronic P-85 and 600 mg of Pluronic L-64 in 50 mL of PBS at 4°C. The mixture is incubated for 30 minutes at 37°C and then sterilely filtered through a 0.22 μ m filter. This is mixed with a solution of 1 mg of sterile lyophilized daunorubicin powder and 5 mg of glucose dissolved in 50 mL of PBS and the mixture is incubated for 30 minutes at 37°C.
- 10 The composition can be stored in the dark at room temperature for 7 days without any essential loss of activity or can be lyophilized and stored for at least 1 year in the dark at room temperature.

Example 21


- 15 A further composition suitable for parental administration is prepared by dissolving 100 mg of sodium ascorbate in a 9% aqueous solution of sodium chloride. To one-half of this solution are added at 4°C 400 mg of Pluronic P-85 and 600 mg of Pluronic L-64. The mixture is incubated for 30 minutes at 37°C and then sterilely filtered through a 0.22 μ m filter. Separately 10 mg of sterile lyophilized daunorubicin powder and 50 mg of glucose are dissolved in the remaining sodium ascorbate-sodium chloride solution and the two solutions are mixed and incubated for 30 minutes at 37°C.
- 20
- 25

This composition can be stored for 30 days in the dark at room temperature without any essential loss of activity or can be lyophilized and stored for at least 1 year in the dark at room temperature.

What is claimed is:

1. In a pharmaceutical composition comprising an anti-neoplastic agent, the improvement in which said agent is incorporated into micelles of at least one block copolymer of poly(oxyethylene)-poly(oxypropylene) in which the ratio of poly(oxypropylene) blocks to the poly(oxyethylene) blocks is from about 0.25 to about 1.5 and the micelles have an average diameter of from about 10 to about 25 nm.

2. A composition according to claim 1 wherein each block copolymer of poly(oxyethylene)-poly(oxypropylene) is independently represented by the formula:



in which each of x and z , independently of the other, has a value of from about 5 to about 100 and y has a value of from about 20 to about 80.

3. A composition according to claim 1 in which the micelles are formed from a plurality of block copolymers of poly(oxyethylene)-poly(oxypropylene) having different hydrophobic/hydrophilic properties.

4. A composition according to claim 3 in which the composite of the block copolymers satisfies the equation:

$$N = \frac{y_1 * a}{(x_1 + z_1) * (a + b)} + \frac{y_2 * b}{(x_2 + z_2) * (a + b)} * 1.32$$

in which:

- y_1 and y_2 are the number of oxypropylene units in the first and second block copolymers, respectively;
 x_1 and z_1 are number of oxyethylene units in the first block copolymer;
5 x_2 and z_2 are number of oxyethylene units in the second block copolymer;
 a is the weight proportion of the first block copolymer; and
 b is the weight proportion of the second block
10 copolymer
such that the value of N is from about 0.25 to about 1.5.
5. A composition according to claim 1 wherein the anti-neoplastic agent is an alkaloid, antibiotic, or antifolate.
- 15 6. A composition according to claim 5 in which the anti-neoplastic agent is selected from the group consisting of vinblastine, colchicine, demecoline, daunorubicin, doxorubicin, mitomycin C, N-methyl mitomycin C, bleomycin A_2 , methotrexate, aminopterin, and dideazatetrahydrofolic
20 acid.
7. A composition according to claim 6 in which the anti-neoplastic agent is daunorubicin.
8. An aqueous dispersion of a quantity of a composition according to claim 1 at least sufficient to provide an
25 effective dose of said anti-neoplastic agent upon parenteral administration.
9. In the method of combatting neoplasms through administration of an anti-neoplastic agent, the improvement comprising administering said anti-neoplastic agent incorporated into micelles of at least one block copolymer of
30 poly(oxyethylene)-poly(oxypropylene) in which the ratio of poly(oxypropylene) blocks to the poly(oxyethylene) blocks is from about 0.25 to about 1.5 and the micelles have an average diameter of from about 10 to about 25 nm.

A. CLASSIFICATION OF SUBJECT MATTER
 IPC 5 A61K9/107 A61K47/10

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
 IPC 5 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US,A,5 039 527 (TABIBI E. ET AL) 13 August 1991	1,2,8,9
Y	see column 2, line 25 - column 3, line 8 see column 4; example 1	5-7
Y	--- DATABASE WPI Section Ch, Week 8443, Derwent Publications Ltd., London, GB; Class B02, AN 84-265868 & JP,A,59 161 313 (MORISHITA PHARM KK) 12 September 1984 see abstract --- -/--	5,6

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

24 January 1994

Date of mailing of the international search report

01.02.94

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
 Fax (+ 31-70) 340-3016

Authorized officer

Boulois, D

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>DATABASE WPI Section Ch, Week 8403, Derwent Publications Ltd., London, GB; Class B02, AN 84-013559 & JP,A,48 088 220 (KYOWA GAS CHEM IND) 19 November 1973 see abstract</p> <p>---</p>	5,6
Y	<p>EP,A,0 219 922 (VESTAR RESEARCH INC) 29 April 1987 see page 6, line 5 - line 35</p> <p>---</p>	7
A	<p>WO,A,88 07365 (RANNEY D.) 6 October 1988 see page 4, line 11 - line 15 see page 6, line 16 - line 24 see page 26, line 21 - line 27</p> <p>---</p>	1
A	<p>EP,A,0 397 307 (RESEARCH DEVELOPMENT CORPORATION OF JAPAN) 14 November 1990 see page 7 - page 8; example 2</p> <p>---</p>	1
A	<p>PHARMAZIE vol. 47, no. 6 , June 1992 , BERLIN pages 439 - 443 XP304546 SHA YANG L. ET AL 'IN VITRO RELEASE, PHARMACOKINETIC AND TISSUE DISTRIBUTION STUDIES OF DOXORUBICIN HYDROCHLORIDE (ADRIAMYCIN HCL) ENCAPSULATED IN LIPIODOLIZED W/O EMULSIONS AND W/O/W MULTIPLE EMULSIONS' see page 442</p> <p>---</p>	1
A	<p>CHEM. PHARM. BULL. vol. 40, no. 8 , August 1992 pages 2224 - 2226 MIYAZAKI S. ET AL 'ANTITUMOR EFFECT OF PLURONIC F127 GEL CONTAINING MITOMYCIN C ON SARCOMA 180 ASCITES TUMOR IN MICE' see page 2224</p> <p>---</p>	1
A	<p>J. OF PARENTERAL SCIENCES AND TECHNOLOGY vol. 41, no. 1 , 1 January 1987 pages 31 - 33 TARR B.D. ET AL 'A NEW PARENTERAL VEHICLE FOR THE ADMINISTRATION OF SOME POORLY WATER SOLUBLE ANTI CANCER DRUGS' see page 31 - page 32</p> <p>-----</p>	1

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A-5039527	13-08-91	NONE	
EP-A-0219922	29-04-87	AU-B- 587232	10-08-89
		AU-A- 5418486	16-04-87
		CA-A- 1272686	14-08-90
		JP-A- 62087514	22-04-87
		US-A- 4769250	06-09-88
WO-A-8807365	06-10-88	US-A- 4925678	15-05-90
		AU-B- 607494	07-03-91
		AU-A- 1627588	02-11-88
		DE-A- 3881881	22-07-93
		EP-A, B 0352295	31-01-90
		JP-T- 4504404	06-08-92
		US-A- 5108759	28-04-92
EP-A-0397307	14-11-90	JP-A- 2300133	12-12-90
		AU-B- 640927	09-09-93
		AU-A- 5469990	15-11-90